

Master your TPN infusion with confidence

PREPARE YOUR SPACE

Workspace

- Clean with disinfecting solution.
- Wash hands with soap & water (or alcohol-based hand sanitizer) for at least 20 sec. Dry with clean paper towel.

TPN Bag

- Remove TPN bag from refrigerator at least 2 hrs. before initiating infusion to allow it to reach room temperature.
- Check label on TPN bag to confirm correct name, date of birth and expiration date.
- Inspect bag for leaks, discoloration or particles. Do not use if you notice any issues and call 1-800-466-3876 for help.

Supplies (gather and arrange supplies as directed by this guide)

- Alcohol pads
- Pre-filled normal saline syringes and Heparin syringes (if prescribed)
- TPN bag and additives (example: multivitamins, insulin, etc.)
- Syringes with needles for additives
- Tubing and Infusion pump

TPN Bag for Administration

- Inject any prescribed additives into the TPN bag (refer to guide for step-by-step instructions).

S

Saline



S

Saline



H

Heparin

(If Ordered)



● 100 units/mL

● 10 units/mL

Remove air from Normal Saline syringe before use.

Use a new alcohol swab for each step. Scrub the catheter's needleless connector with swab for at least 15 sec.

Flush catheter using pulsatile flush technique.

After flushing and locking the catheter, apply a disinfection cap to the needleless connector.